

CS 530 - Fall 2018
Homework 1

Due: Tuesday, September 18 - submit via Gradescope

Reading : Section 4.2, pages 75-82 and Sections 28.1 and 28.2 pages 813-830 of the textbook

Problems: Please limit your answer to the following problems to at most 2 pages each.

1. Strassen's method

Assume you are handed an algorithm which multiplies two 2 by 2 matrices and uses 7 integer multiplications and 18 integer additions to do this.

i. Show how to use this algorithm iteratively to multiply two 4 by 4 matrices. Calculate exactly how many integer multiplications and integer additions are done? You should show your work and should compare the answer you get to the number of multiplications and adds which would be done if you turn the 4 by 4 algorithm into block multiplication of 2 by 2 matrices.

ii. In Section 4.2 of the textbook it is shown that Strassen's method uses only $O(n^{\log_2 7})$ multiplications when n is an even power of 2. (You need not prove this.)

Using this result describe how you can get the same $O(n^{\log_2 7})$ multiplications even if you start with n by n matrices with n not an even power of 2.

2. A permutation matrix P is an n by n Boolean matrix with exactly one 1 in each row and one 1 in each column.

P is called a permutation matrix because if you multiply P by any n by 1 column vector V then PV is a permutation of V .

Given a permutation matrix P :

i. Explain what permutation of $1, 2, \dots, n$ the matrix P corresponds to. Is P invertible? If so, explain how to construct the inverse of P .

ii. What are the possible values of the determinant of P ? What is the relationship between the determinant of P and $\text{sign}(Q)$, where Q is the permutation which corresponds to the matrix P ? Here $\text{sign}(Q)$ is the number of flips (i.e., interchanges of permuted integers) that need to be done to go from the identity permutation to Q . Explain why the relationship holds.

iii. Let $P =$

$$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

Construct the inverse of P and compute P 's determinant.

iv. Give an example of a 3 by 3 permutation P which has no LU decomposition.

3. i). Prove that for any n greater than or equal to 2 there is a non-singular square (n by n) matrix which has no LU decomposition with L unit lower triangular and U upper triangular.

Give a specific example of your matrix when $n=3$ and show its LUP decomposition.

(ii). Give an example of a singular 2 by 2 matrix A and a singular 3 by 3 matrix B which have an LU decomposition. Both A and B should be non-zero matrices and you should write the L and the U for both of them.